



Marker-assisted Selection in Fish: A Review

F. Eze^{1*}

¹*Department of Fisheries and Aquaculture, Faculty of Marine Environmental Management, Nigeria Maritime University, Okerenkoko, Warri South, Delta State, Nigeria.*

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

DOI: 10.9734/AJFAR/2019/v3i430038

Editor(s):

(1) Dr. Jorge Castro Mejia, Department of El Hombre Y Su Ambiente, Universidad Autonoma Metropolitana Xochimilco, Mexico.

Reviewers:

(1) K. D. Mini, Mahatma Gandhi University, India.

(2) Babatunde Akeem Saka, University of Ibadan, Nigeria.

(3) Theodoros Mavraganis, Holar University College, Iceland.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50277>

Received 12 May 2019

Accepted 22 July 2019

Published 30 July 2019

Review Article

ABSTRACT

The important economical traits like body growth, resistance to diseases, meat quality, etc. highly influence the profitability of food animals including fishes. The main target of every selective breeding programme is to produce improved traits offspring's. However, improvement of performance traits through traditional phenotype-based selection needs several generations to optimise these characters. Marker-Assisted Selection (MAS) is a type of indirect method of selection of better performing breeding individuals. MAS is beneficial when the traits are difficult, expensive to measure and has both low heritability and recessive traits. MAS facilitates the exploitation of existing genetic diversity in breeding populations and can be used to improve desirable traits in livestock. MAS depends on identifying the link between a genetic marker and Quantitative Traits Loci (QTL). The distance between marker and target traits determines the association of the marker with the QTL. After identifying the markers linked to QTL, they can be used in the selective breeding programme to select the brooders having better genetic potential for the targeted trait. Improvement of performance traits through MAS is fast and more accurate and allows us to understand the genetic mechanism affecting performance traits.

Keywords: *Marker-assisted selection; quantitative traits loci; genetic diversity; trait.*

1. INTRODUCTION

Marker-Assisted Selection (MAS) is a type of biotechnology that uses molecular genetic markers as a criterion for selecting desired traits Ashraf et al. [1] Marker Assisted Selection (MAS) is an indirect selection process where a trait of interest is selected not based on the trait itself but on a marker linked to it Ribaut and Ragot [2].

MAS is considered a “revolutionary” approach to traditional tree breeding as it allows breeders to select individuals based on their genotypes, rather than being restricted to phenotypic characteristics (Boopathi et al., 2013).

Sax [3] was the first to show how genetic factors influencing quantitative traits can be identified using markers.

Recently MAS became a very popular method of indirect selection for production of the genetically improved in aquaculture breeding programme offspring's. Most of the performance traits including growth or disease resistance are controlled by multiple genes and are therefore inherited as quantitative traits, analysis of their associated quantitative trait loci (QTL) is an essential part of aquaculture genomics (Liu and Cordes, 2004). QTLs are largely unknown genes that affect performance traits (such as growth rate and disease resistance) and these are important to breeders.

MAS in a breeding context involves scoring indirectly for the presence or absence of a desired phenotype or phenotypic component based on the sequences or banding patterns of molecular markers located in or near the genes controlling the phenotype. The sequence polymorphism or banding pattern of the molecular marker is indicative of the presence or absence of a specific gene or chromosomal segment that is known to carry a desired allele Brumlop and Finckh [4].

Marker-assisted selection method (MAS) or genome-wide marker-assisted selection method (G-MAS) was not widely used in aquaculture, but nowadays its use is increasing due to its ease of use and quicker than traditional phenotype-based selection. Now it becomes a fertile field of research for the aquaculture researchers to discover novel genetic marker that can be used to link with the QTLs in selective breeding programmes (Hauser et al., 2011; Dichmont et al., 2012; Abdul-Muneer, 2014).

In order to manage individual species effectively, identification of different species from a mixed catch becomes important. DNA markers are widely being accepted not only to obtain information about gene flow and allele frequencies in aquaculture practices but also to identify hybrids. The majority of the markers, which are used in inter- and intra-specific disparity, include Random Amplified Polymorphic DNA (RAPD) for species and sub-species identification done in tilapia (Bardakci and Skibinski [5], and iso-enzyme used in intraspecific variations in Sparidae species (Alarcón and Alvarez [6]. Similarly, Nijman et al. (2003) reported the use of mtDNA markers as an important tool in rapid detection of hybridization between species and subspecies of livestock.

Markers tend not to have any biological effect, but rather can be thought of as notable and constant points of reference within the genome (Guimaraes, et al., 2007). Markers can be found within the desired gene or, more commonly, linked to a gene determining a trait of interest (Brumlop and Finckh, 2011; Guimaraes et al., 2007). Unlike genetic engineering, MAS does not alter the original DNA Vogel and Van Aken [7]; instead it uses genetic marker to identify naturally-occurring genetic variations among individuals, with the intent of selecting those with the best potential to meet desired criteria and objectives.

Marker Assisted Selection (MAS) provides several other benefits to breeders, in that it can select for genes that demonstrate low heritability, have recessive alleles, and are difficult, expensive, or time exhaustive to determine phenotypically Boopathi [8], Brumlop and Finckh [4], Xu et al. [9]. MAS also allows for gene pyramiding or combining multiple genes within the same breeding line, while having fewer unintentional losses and fewer selection cycles Boopathi [8], Xu and Crouch [9].

Furthermore, MAS may be viewed by the public with more support than genetic engineering as breeders are not manually manipulating the genes, and thus all offspring inheritance occurs naturally Vogel and Van Aken [7]. It is also believed that genetic markers may be important in the assessment, conservation and use of diversity in germplasm and varieties Brumlop and Finckh [4].

Molecular marker maps have been constructed for a number of aquaculture species, e.g. tilapia,

Clarias, giant tiger prawn, kuruma prawn, Japanese flounder and Atlantic salmon, although their density is generally low Nichols et al. [10]. As many preferred traits are not observed until maturity, MAS eliminates this waiting period by allowing for the early selection of desired genotypes at the seedling stage Yanchuk [11].

The desirable phenotypic variations in the performance traits of fishes are used to increase the aquacultural yield, improve incomes of farmers and enhances food security through selective breeding by choosing better-performed individuals. However, phenotype-based selection needed considerable time to optimise the traits, so researchers are now moving from phenotype based selection to genotype-based selection. The absence of a molecular marker is the main limiting factor for the realization of genotype based selection potentials in fishes. However, with the advent of DNA-based genetic markers in the late 1970s and now the ease of the marker discovery through the next generation sequencing allowed researchers to identify large numbers of markers spreads throughout the genome of any species of interest. The markers are used to detect linkage with the traits of interest, thus allowing MAS finally to become a reality Peterson et al. [12]. This paper aims to provide information regarding the technical aspect of MAS and the current application in fisheries and Aquaculture in other to increase high quality production within a period of time.

2. MARKER ASSISTED SELECTION

Incorporation of marker information into breeding programs in aiding identification and selection of superior individuals has been widely studied Bernardo [13]; Han et al., 1997; Xie and Xu [14], Romagosa et al., 1999; Ayoub et al., 2003; Jordan et al., 2003).

Molecular markers in aquaculture and fisheries have been used for over 50 years Ryman and Utter [15]; Liu and Cordes, 2004) and their use has steadily increased over the last two decades (Park and Moran, 1994; Chauhan and Rajiv, 2010; Dichmont et al., 2012; Abdul-Muneer, 2014).

An important factor in MAS is the accuracy of estimating the genetic effects related to the trait of interest. In contrast to genetic engineering (GE), MAS does not alter the original DNA. Rather, it identifies whether the desired trait(s) are being expressed, so that individuals with the

best potential can be selected Andersson et al. [16].

Molecular marker analysis allows the identification of genome segments, so called Quantitative Trait Loci (QTL), contributing to the genetic variance of a quantitative trait and thus to select superior genotypes as these loci Cannai et al. [17]. Allelic variation in genetic markers can be linked to the variation in traits of economic interest, and thus the marker provides DNA level information on the inheritance of the traits.

The practical use of markers in selection can be roughly divided into three classes:

- 1) Removing genetic disorders,
- 2) Marker breeding value-selection, and
- 3) Genomic selection.

2.1 MAS versus Phenotypic Selection

Marker-Assisted Selection (MAS) will probably never replace Phenotypic Selection (PS) entirely. There is no general pattern by which it can be predicted whether MAS or PS will be more useful. Empirical comparisons of MAS and PS for increasing gain from selection have been made in several studies. The outcomes of these studies are conflicting. In some studies, MAS is reported to be more effective/efficient than PS (Yousef and Juvik 2001; Abalo et al., 2009) while other studies considered the two methods equal (Van Berloo and Stam 1999; Willcox et al.[18], Hoeck et al., 2003; Moreau et al. [19]. In a third group of studies PS proved to be more effective/efficient than MAS (Davies et al. [20], Wilde et al., 2007) and in other comparisons the effectiveness/efficiency of MAS and PS varied within the same study, depending on the populations or on the trait selected for Flint Garcia *et al.*, [21]; Robbins and Staub 2009).

2.2 Limitations of MAS

- Cost
- Requirement of technical skill
- Automated techniques for maximum benefit

2.3 Advantages of MAS

In addition to the cost and time savings described above, for a number of breeding scenarios, MAS methods are likely to offer significant advantages compared with

conventional selection methods. These scenarios assume the availability of markers for multiple traits and take into consideration the advantages of MAS under optimum situations Dreher et al. [22], Dudley [23].

1. Gene stacking for a single trait: MAS offers potential savings compared with conventional selection when it allows breeders to identify the presence of multiple genes/alleles related to a single trait, and the alleles do not exert individually detectable effects on the expression of the trait.
2. Early detection: MAS offers potential savings compared with conventional selection when it allows alleles for desirable traits to be detected early, well before the trait is expressed and can be detected phenotypically. This benefit can be particularly important in species that grow slowly.
3. Heritability of traits: Up to a point, gains from MAS increase with decreasing heritability. However, due to the difficulties encountered in QTL detection, the gains are likely to decline beyond a certain threshold heritability estimate.

2.4 Disadvantages of MAS

Perhaps the greatest disadvantage of MAS is the time and financial investment required to develop markers that are widely applicable for traits of agronomic importance. Often a marker developed in one or a few related genotypes will not work for other genotypes in a breeding scheme due to allelic effects. Furthermore, development of markers, particularly for QTL, is complicated by epistatic interactions and the critical need for good quality phenotypic data.

2.5 Quantitative Trait Loci

In fish, several QTL studies have been published; in salmonids Jackson *et al.* [24]; Johansen 1999; Robinson *et al.* [25], Sakamoto *et al.*[26], Marfyniuk 2001, Ozaki *et al.* [27] Somorger 2001. Tao and Baiding 2003), in catfish Liu *et al.* [28], in tilapia Cnaani *et al.* [17] and in silver barb Hussain *et al.* 2002.

Marker Assisted Selection (MAS) is followed by two steps, detection of molecular markers associated with quantitative trait locus (QTL) and application of those markers.

The position of the chromosome that controls the economical important trait is termed as QTL.

The concepts for detecting QTL were developed more than 90 years ago Sax [3]. In aquaculture species, much effort has been applied for QTL mapping. QTLs are mapped by linkage disequilibrium with molecular markers exhibiting Mendelian segregation. Economically important traits are controlled by the single or group of gene.

The basic concept of QTL studies is to know the number and location of loci associated with phenotypic traits Mackay [29]; Mauric io, 2001; Burt and Hocking, 2002; Erickson *et al.*, 2004). Thus, candidate gene or molecular markers, resulted by QTL mapping, could be used in MAS Groen *et al.*, [30]. QTL detection is an ongoing effort in aquaculture species. More than 37 important traits have been located in about 20 aquaculture species.

QTL mapping is the practical application of marker-assisted selection in aquaculture (Al-Samarai, 2015). With rapid advancement of molecular technology, it is now possible to use molecular marker information to map major quantitative trait loci (QTLs) on chromosomes Paterson *et al.* [31], Hilbert *et al.* 1991, Jacob *et al.* [32], Stuber *et al.*, 1992. Once QTL for a trait are identified, individuals can be selected for breeding on the basis of marker alleles that segregate with favourable phenotypes (Lande and Thompson, 1990). This strategy, known as marker-assisted selection (MAS), is particularly useful for traits that cannot be measured on selection candidates directly, notably disease resistance or meat quality traits Sonesson [33].

2.5.1 QTL detection in fish

A number of genetic maps have been developed specifically to locate QTL in several fish species. The first of such map was produced in Zebrafish insert scientific name Postlethwait *et al.*,[34]; Shimoda *et al.*[35], which is a non-aquacultural species. Among cultivable fish groups low-density maps have been developed for salmonids Sakamoto *et al.*[36]; Ghabi 2001) for catfish (Liu *et al.*, 2003; Poopuang and Na-Nakorn 2004) for tilapia Kocher *et al.*[37]; Cnaani *et al.*[17], for Japanese flounder (Sanchez *et al.*, 2003), for red sea beam (Sakamoto *et al.*, 2003), for Oyster (Yu and Geso 2003), and for shrimp ([Http://shrimppmap.tag.csiro.au](http://shrimppmap.tag.csiro.au)).

2.5.2 QTL mapping in fish

Although in fish several studies have confirmed the existence of significant genetic variation for quantitative traits at commercial importance Kaue *et al.* [38] and have recognized the potential of MAS for their genetic improvement (Flint and Mott, 2001). Thus far, very few QTL for production traits have been identified in fish Sonesson [39]. Much effort is devoted to QTL mapping for growth, feed conversion efficiencies, disease resistance, fecundity, and spawning time (Dunham *et al.*, 2001).

Several QTL studies have been published in rainbow trout for temperature tolerance (Jackson *et al.*, 1998). Danzmann *et al.*, 1998, Perry [40], spawning time Sakamoto *et al.* [41]; fish back *et al.* 2000, O' Malley 2001); growth (Martynick 2001), disease resistance (Ozaki *et al.*, 2001), and fitness traits Somorjai [42]. Other notable QTL studies published in aquaculture fish species include: in tilapia for temperature and salinity tolerance (Streadman and Kocher 2002; Cnaan *et al.*, 2003), in catfish for feed conversion efficiency and bacterial septicaemia resistance (Liu 2003), in guppy for growth (Nakajima and

Table 1. QTL studies in selected aquaculture species

Species	Traits	References
Arctic charr	Body weight and sexual maturation; Salinity tolerance	Küttner <i>et al.</i> , 2011
Asian seabass	Resistance against viral nervous necrosis disease Growth-related trait Omega-3 fatty acids	Wang <i>et al.</i> [47] Xia <i>et al.</i> , 2014
Atlantic salmon	Growth traits and flesh colour Resistance against IPN Late sexual maturation	Baranski <i>et al.</i> , 2010; Tsai <i>et al.</i> , 2014; Moen <i>et al.</i> , 2009 ; Houston <i>et al.</i> , 2008; 2010 Gutierrez <i>et al.</i> , 2014
Catfish	Columnaris disease resistance ESC disease resistance Hypoxia tolerance Heat stress Head size	Geng <i>et al.</i> , 2015 Wang <i>et al.</i> , 2013; Zhou <i>et al.</i> , 2017 Wang <i>et al.</i> , 2016; Jin <i>et al.</i> , 2016 Geng <i>et al.</i> , 2016
Common carp	Common carp Morphometric traits Swimming ability	Zhang <i>et al.</i> , 2011 Boulton <i>et al.</i> , 2011 Laghari <i>et al.</i> , 2014
Eastern oyster	Disease resistance	Yu and Guo [48]
European seabass	Growth, body weight Morphometric traits and stress Response	Louro <i>et al.</i> , 2016 Massault <i>et al.</i> , 2010
Pacific white shrimp	Growth parameters	Andriantahina <i>et al.</i> , 2013
Giant tiger prawn	Disease resistance and sex determination	Robinson <i>et al.</i> , 2014
Japanese flounder	<i>Vibrio anguillarum</i> resistance	Wang <i>et al.</i> , 2014
Pacific oyster	Growth Resistance against summer mortality Viability	Guo <i>et al.</i> , [49] Sauvage <i>et al.</i> , 2010 Plough and Hedgecock, 2011 Plough <i>et al.</i> , 2016
Gilthead seabream	Skeletal deformities Sex determination and body growth	Negrín-Báez <i>et al.</i> , 2015 Loukovitis <i>et al.</i> , [50] Massault <i>et al.</i> , 2011
Rainbow trout	Growth related traits	Kocmarek <i>et al.</i> , 2015; Wringe <i>et al.</i> [51]; Leder <i>et al.</i> , 2006.
Tilapia	Growth traits Sex	Liu <i>et al.</i> , [52] Wang <i>et al.</i> , 2015 Palaikostas <i>et al.</i> , 2015

Taniguchi 2002), in at fautic salmon for infectious anaemia resistance (Moen et al., 2003 and in Artic charr for growth rates and fitness traits (Johansen 1999, Somorjai 2001).

In salmonids, QTL have been found related to body weight and size Martyniuk et al.[43]; O'Malley et al.[44]; Reid et al. [45], for colouration pattern (Streelman, Albertson and Kocher, 2003) and for one form of albinism (Nakamura et al., 2001). Zimmerman et al. [46] found QTL for pyloric caeca number, a trait related to feed conversion efficiency.

2.5.3 QTL analysis

2.5.3.1 QTL for growth traits

Growth is one of the most important economic traits of all aquaculture species. Up to 2012, QTL analyses have been conducted in more than 20 aquatic species Yue et al. [53] and growth was the most popular trait studied. Wang et al., [54] used 380 F1 Asian seabass to identify five major QTLs and 27 potential QTLs. Of them, three major QTLs for body weight, length, and body length were located at a similar linkage group 2 (LG2) position with the nearby Lca287 microsatellite and accounted for 28.8%, 58.9%, and 59.7% of the phenotypic variations. The other two major QTLs for body weight were located at another LG2 position. These five major QTLs have been confirmed in two other Asian seabass populations Wang et al. [55]. Further QTL fine mapping of the Asian seabass growth trait identified three candidates "growth genes" (CATHEPSIN D, KCTD15, and CSMD2) affecting body weight, body length, and total length (Wang et al., 2011). The function of the *cathepsin D* gene in humans involves cell proliferation and cell growth; therefore, *cathepsin D* may also be a major "growth gene" in Asian seabass. O'Malley et al., (O'Maller et al., 2008) identified QTLs for body weight in rainbow trout on 10 different LGs.

Wringe et al. (2010) used additional backcrossed families and SSR markers to confirm the O'Malley et al. (2003),s results and found several major candidate growth genes (e.g., *GH2* and *Pax7*). Reid et al., 2004 identified a QTL for body weight in two LGs (AS8 and 11) of Atlantic salmon, and reported that it was homologous to the growth QTL in rainbow trout. Houston et al. [56,57] identified QTLs for body weight in Linkage group 1 (LG1) and LG5 of Atlantic salmon. Gutierrez et al., (2012) further used a

6.5 K Single Nucleotide Polymorphisms (SNP) chip to identify QTLs in six LGs at the genomic level. Cnaani et al. (2004) identified a QTL for tilapia growth on LG23, which is the linkage group with the genetic sex-determining region. Song et al.[58,59] used 1487 SSRs to produce a high-density genetic linkage map and successfully identified a QTL affecting body weight in LG14 of Japanese flounder.

Some reports have used a candidate gene approach to identify growth-related genes and molecular markers in fish. Tao and Boulding (2003) found polymorphisms in the growth hormone gene (*GH*) that were significantly associated with growth rate of Arctic charr (*Salvelinus alpinus*). Li et al. [60] reported an SNP in the insulin-like growth factor- (IGF)1gene 5' untranslated region (UTR) of largemouth bass (*Micropterus salmoides*). Sun et al. (2012) reported that two SNPs in exon 3 of the myostatin (*MSTN*) gene were significantly related to body weight and Fulton's factor in common carp. Liu et al. (2012) also found that a SNP in the *MSTN* 3' UTR was very significantly associated with total length, body length, and body weight of bighead carp.

2.5.3.2 QTL for feed conversion rate

Food Conversion Ratio (FCR) is one of the most important economic traits in fish, as fish with a better FCR increase profits.

Liu (2005) used Amplified Fragment Length polymorphisms (AFLP) markers to construct a catfish genetic map and found a QTL associated with FCR. Zimmerman et al. [46] revealed three QTLs for the number of pyloric caeca in three LGs of rainbow trout, and this is an important index associated with FCR.

Food Conversion Ratio (FCR) studies have also been reported in common carp from the Heilongjiang Fisheries Research Institute of the Chinese Academy of Fishery Sciences Wang et al. [61].

2.5.3.3 QTL for sex determination

Sex phenotype and sex determination in fish have specific evolutionary status and diversity. Males and females of some species have significant differences in growth rate or commercial value; therefore, mono sex fish culture is a promising strategy. The sex-determining (SD) loci and QTLs have been

studied in a limited number of fish, such as tilapia (Lee et al., 2004) rainbow trout (Alfaqih et al., 2009) and salmonids (Davidson et al. [62]. Previous studies have demonstrated that sex QTLs are located on LG1, 2, 3, 6, and 23 of tilapia (Cnaani et al., 2004; Lee et al., 2004; Cnaani et al., 2008) Eshel et al. [63] reported a major candidate sex QTL that is considered the sex determining region in tilapia. Fifty-one genes in this region have been annotated, and 10 have been confirmed.

The anti-Müllerian hormone gene is the most differentially expressed gene in male and female tilapia. Sun et al., (2014) recently published several sex-specific markers, and one is tightly linked with the sex-determining region discovered by Eshel et al., [63]. The sex-determining locus in rainbow trout is located on the LG of RT10, and this locus also significantly affects thermo-resistance and body length. The sex-determining regions in Artic charr Moghadam et al. [64] brown trout Gharbi et al.[65] and Atlantic salmon Gilbey et al. [66] are located on the LGs of AC4, BT28, and AS1, respectively.

Woram et al.,[67] compared LGs of sex-determining loci in four salmonids and found that although the nucleotide sequences flanking the sex-determining loci were well-conserved, the SD LGs were diverse, suggesting that the regions underwent different recombination events.

Loukovitis et al. [68] located growth and sex-determining QTLs in gilthead sea bream and showed that these two traits have similar genetic control in LG21. Martínez et al., [69] located a sex QTL on LG5 of turbot and proposed a ZZ/ZW sex-determining mechanism. Viñas et al., (2012) also found a major sex QTL on turbot LG5. These findings suggest that the sex-determining genes may occur on turbot LG5. Song et al. [58,59] used high-density genetic maps to locate seven sex QTLs on the half-smooth tongue sole LG1f, LG14f, and LG1m.

Additional study by Chen et al. [70] provided insight into ZW sex chromosome evolution and identified sex-determining genes, such as *dmrt1* and *neurl3*.

2.5.4 Factors affecting QTL analyses

The power of mapping QTL can be influenced by a number of factors, such as genetic properties of QTL, experimental design, environmental

effects, marker density and informativeness, genotyping errors and precision of trait measurement. Details about how these factors influence the power of QTL mapping can be found in some very good reviews (Crosses 2001; Flint and Mott 2001; Doerge 2002).

2.5.5 Methods of detecting QTL

Basically, three methods are frequently used for mapping QTL and estimating their effects, namely Single-Marker Association Analysis (SMAA), Simple Interval Mapping (SIM) and Composite Interval Mapping (CIM) (Crosses 2001; Flint and Mott 2001; Doerge 2002).

2.6 Current Status of Applications of MAS in Fish

Molecular marker maps have been constructed for a number of aquaculture species, e.g. tilapia, catfish, giant tiger prawn, kuruma prawn, Japanese flounder and Atlantic salmon, although their density is generally low. Density is high for the rainbow trout, where the map published in 2003 has over 1300 markers spread throughout the genome – the vast majority are AFLPs but it also includes over 200 microsatellite markers (Nichols et al., 2003). Some QTLs of interest have been detected (e.g. for cold and salinity tolerance in tilapia and for specific diseases in rainbow trout and salmon).

3. CONCLUSION

In a recent review of MAS in fish breeding schemes, Sonesson (2003) suggested that MAS would be especially valuable for traits that are impossible to record on the candidates for selection such as disease resistance, fillet quality, feed efficiency and sexual maturation, and concluded that MAS is not used in fish breeding schemes today and that the lack of dense molecular maps is the limiting factor. Marker Assisted Selection (MAS) has become a valuable tool in selecting organisms for desirable traits. MAS is expected to increase genetic gain compared to traditional breeding programs and reduce the cost of progeny testing by early selection. The application of MAS in breeding programmes depends on the knowledge of breeders about variable marker information.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Ashraf M, Akram NA, Foolad MR. (Eds.). Marker-assisted selection in plant breeding for salinity tolerance. 2012;913:305–333. DOI:10.1007/978-1-61779-986-0.
2. Ribaut JM, Ragot M. Marker-assisted selection to improve drought adaptation in maize: The backcross approach, perspectives, limitations and alternatives. *J Exp Bot.* 2007;58:351-360.
3. Sax K. The association of size differences with seed-coat pattern and pigmentation in *Phaseolus vulgaris*. *Genetics.* 1923;8:522-560.
4. Brumlop S, Finckh MR. Applications and potentials of marker assisted selection (MAS) in plant breeding. 2011;178. Available: http://www.bfn.de/0502_skripten.html
5. Bardakci F, Skibinski DOF. Application of the RAPD technique in tilapia fish species and subspecies identification. 1994;73:117–123. DOI:10.1038/hdy.1994.110.
6. Alarcon JA, Alvarez MC. Genetic identification *Sparidae* species by isozyme markers. Applications to Interspecific Hybrids. *Aquaculture.* 1999;173:95-103.
7. Vogel B, Van Aken J. Smart breeding - Marker-assisted selection: A non-invasive biotechnology alternative to genetic engineering of plant varieties Amsterdam, the Netherlands. 2009;28.
8. Boopathi NM. Marker-assisted selection. In genetic mapping and marker assisted selection: Basics, practice and benefits. 2013a;173–186. DOI:10.1007/978-81-322-0958-4
9. Xu Y, Beachell H, McCouch SR. A marker based approach to broadening the genetic base of rice in the USA. *Crop Sci.* 2008;44:1947–1959.
10. Nichols KM, Bartholomew J, Thorgaard GH. Mapping multiple genetic loci associated with *Ceratomyxa shasta* resistance in *Oncorhynchus mykiss*. *Dis. Aquat. Org.* 2003;56:145–154.
11. Yanchuk AD. The role and implications of biotechnology in forestry. *Food and agriculture organization of the United Nations, Unasyuva.* 2002;(30):18–22.
12. Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. Double digest RADseq: An inexpensive method for de novo SNP discovery and genotyping in model and non-model species. *PLoS One.* 1990;7:e37135.
13. Bernardo R. Prediction of maize single-cross performance using RFLPs and Biotechnology in Agriculture and Food; 1994. Available: <http://tandfonline.com/doi/book/10.1081/EEBAF>.
14. Xie C, Xu X. Efficiency of multistage marker-assisted selection in the improvement of multiple quantitative traits. *Heredity.* 1998;80:489-498.
15. Ryman N, Utter F. Population genetics and fishery management University of Washington Press, Seattle. 1987;420.
16. Anderson JL, Marí AR, Braasch I, Amores A, Hohenlohe P, Batzel P. Multiple sex-associated regions and a putative sex chromosome in zebrafish revealed by RAPD mapping and population genomics. 2001;3:427-437.
17. Cnaani A, Zilberman N, Tinman S, Hulata G, Ron M. Genome-scan analysis for quantitative trait loci in an F-2 tilapia hybrid. *Mol Genet Genomics.* 2003;272(2):162-172.
18. Willcox MC, Khairallah M, Bergvinson D, Crossa J, Deutsch JA, Edmeades GO, Gonzalez-de-Leon D, Jiang C, Jewell DC, Mihm JA, Williams WP, Hoisington DA. Selection for resistance to southwestern corn borer using marker assisted and conventional backcrossing. *Crop Sci.* 2002;42:1516–1528.
19. Moreau L, Lamarie S, Charcosset A, Gallais A. Economic efficiency of one molecular marker assisted selection for the malt quality traits in barley. *Mol Breeding.* 2000;3:427-437.
20. Davies J, Berzonsky W, Leach G. A comparison of marker-assisted and phenotypic selection for high grain protein content in spring wheat. 2006;152:117-134.
21. Flint-Garcia SA, Darrach LL, McMullen MD, Hibbard BE. Phenotypic versus genomic selection in an established commercial layer breeding program. *Genetics Selection Evolution.* 2003;45:29.
22. Dreher K, Khairallah M, Jean-Marcel R, Monis M. Money matters (I): Costs of field and laboratory procedures associated with conventional and marker assisted maize breeding at CIMMYT. *Molecular Breeding* 2002;11:221-234.
23. Dudley JW. Molecular markers in plant improvement: Manipulation of genes

- affecting quantitative traits. *Crop Sci.* 1993;33:660–668.
24. Jackson TR, Ferguson MM, Danzmann RG, Fishback AG, Ihssen PE, O'Connell M, Crease TJ. Identification of two QTL influencing upper temperature tolerance in three rainbow trout (*Oncorhynchus mykiss*) half-sib families. *Heredity.* 1998;80:143–151.
 25. Robison BD, Wheeler PA, Sundin K, Sikka P, Thorgaard GH. Composite interval mapping reveals a major locus influencing embryonic development rate in rainbow trout (*Oncorhynchus mykiss*). *J. Heredity.* 2001;92:16–22.
 26. Sakamoto T, Danzmann RG, Okamoto N, Ferguson MM, Ihssen PE. Linkage analysis of quantitative trait loci associated with spawning time in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture.* 1999;173:33–43.
 27. Ozaki A, Sakamoto T, Khoo S, Nakamura K, Coimbra MR, Akutsu T, Okamoto N. Quantitative Trait Loci (QTL) associated with resistance/susceptibility to infectious pancreatic necrosis virus (IPNV) in rainbow trout (*Oncorhynchus mykiss*). *Mol. Genet. Genomics.* 2001;265:23–31.
 28. Liu Z, Karsi A, Li P, Cao D, Dunham R. An AFLP-based genetic linkage map of channel catfish (*Ictalurus punctatus*) constructed by using an interspecific hybrid resource family. *Genetics* 2003;165:687–694.
 29. Mackay TFC. The genetic architecture of quantitative traits. *Annu Rev Genet.* 2001;35:303-339.
 30. Groen AF, Crooijmans RPMA, Van Kampen AJA, Van der Beek S, Van der Poel JJ, Groenen MAM. Microsatellite polymorphism in commercial broiler and layer lines. *Proc 5th World Congr Genet Appl Livestock Prod.* 2000;21:95-98.
 31. Paterson A, Lander SE, Hevit JD, Peterson S, Lincoln SE, Lanksley SD. Resolution of quantitative traits into mendelian factors by using a complete linkage map of Restriction Fragment Length Polymorphisms. *Nature.* 1988;325:721-726.
 32. Jacob HJ, Lindpainter K, Lincoln SE, Kusumi RK, Mao YP, Ganten D, Dzau VJ, Lander ES. Genetic mapping of a gene causing hypersensitive rat. *Cell.* 1991;67:213-224.
 33. Sonesson A. Within-family marker-assisted selection for aquaculture species. *Genet Sel Evol.* 2007;39:301–18.
 34. Postlethwait JH, Johnson SL, Midson CN, Talbot WS, Gates M, Ballinger EW, Africa D, Andrews R, Carl T, Eisen JS. A genetic linkage map for the zebrafish. *Science.* 1994;264:699-703.
 35. Shimoda N, Knapik EW, Ziniti J, Sim C, Yamada E, Kaplan S, Jackson D, de Sauvage F, Jacob H, Fishman MC. Zebrafish genetic map with 2000 microsatellite markers. *Genomics.* 1999;58: 219-232.
 36. Sakamoto T, Danzmann RG, Gharbi K, Howard P, Ozaki A, Khoo SK. A microsatellite linkage map of rainbow trout (*Oncorhynchus mykiss*) characterized by large sex-specific differences in recombination rates. *Genetics.* 2000;155:1331-1345.
 37. Kocher TD, Lee WJ, Sobolewska H, Penman D, McAndrew B. A genetic linkage map of a cichlid fish, the tilapia (*Oreochromis niloticus*). *Genetics.* 1998;148:1225–1232.
 38. Kause A, Paananen T, Ritola O, Koskinen H. Direct and indirect selection of visceral lipid weight, fillet weight, and fillet percentage in a rainbow trout breeding program. *Journal of Anim Sci.* 2003;85:3218-3227.
 39. Sonesson AK. A combination of walk-back and optimum contribution selection for fish – a simulation study. *Genet. Sel. Evol.* 2003;37:587–599.
 40. Perry GM, Danzmann RG, Ferguson MM, Gibson JP. Quantitative trait loci for upper thermal tolerance in outbred strains of rainbow trout (*Oncorhynchus mykiss*). *Heredity.* 2001;86:333– 341.
 41. Sakamoto T, Danzmann RG, Okamoto N, Ferguson MM, Ihssen PE. Linkage analysis of quantitative trait loci associated with spawning time in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture.* 1999;173:33–43.
 42. Somorjai IM, Danzmann RG, Ferguson MM. Distribution of temperature tolerance quantitative trait loci in Arctic charr (*Salvelinus alpinus*) and inferred homologies in rainbow trout (*Oncorhynchus mykiss*). *Genetics.* 2001;165:1443–1456.
 43. Martyniuk CJ, Perry GML, Mogahadam HK, Ferguson MM, Danzmann RG. The genetic architecture of correlations among

- growth related traits and male age at maturation in rainbow trout. *Journal of Fish Biol.* 2003;63:746–764.
44. O'Malley KG, Sakamoto T, Danzmann RG, Ferguson MM. Quantitative trait loci for spawning date and body weight in rainbow trout: Testing for conserved effects across ancestrally duplicated chromosomes. *J. Hered.* 2003;94:273–84.
 45. Reid DP, Szanto A, Glebe B, Danzmann RG, Ferguson MM. QTL for body weight and condition factor in Atlantic salmon (*Salmo salar*): comparative analysis with rainbow trout (*Oncorhynchus mykiss*) and Arctic charr (*Salvelinus alpinus*). *Heredity.* 2005;94(2):166-172.
 46. Zimmerman AM, Wheeler PA, Ristow SS, Thorgaard GH. Composite interval mapping reveals three QTL associated with pyloric caeca number in rainbow trout, *Oncorhynchus mykiss*. *Aquaculture.* 2005;247:85–95.
 47. Wang CM, Lo LC, Zhu ZY, Yue GH. A genome scan for quantitative trait loci affecting growth-related traits in an F1 family of Asian seabass (*Lates calcarifer*). *BMC Genomics;* 2006.
 48. Yu Z, Guo X. Genetic linkage map of the eastern oyster (*Crassostrea virginica*) Gmelin. *Biol. Bull.* 2003;204:327–338.
 49. Guo X, Hershberger WK, Cooper K, Chew KK. Artificial gynogenesis with; 2012.
 50. Loukovitis D, Sarropoulou E, Tsigenopoulos CS, Batargias C, Magoulas A, Apostolidis AP, Chatziplis D, Kotoulas G. Quantitative trait loci involved in sex determination and body growth in the gilthead sea bream (*Sparus aurata* L.) through targeted genome scan. *PLoS ONE.* 2011;6:e16599.
 51. Wringe B, Devlin R, Ferguson M, Moghadam H, Sakhrani D, Danzmann R. Growth-related quantitative trait loci in domestic and wild rainbow trout (*Oncorhynchus mykiss*). *BMC Genetics.* 2010;11(63).
 52. Liu F, Sun F, Xia JH, Li J, Fu GH, Lin G. A genome scan revealed significant associations of growth traits with a major QTL and GHR2 in tilapia. *Sci Rep.* 2014;4:7256.
 53. Yue GH. Recent advances of genome mapping and marker-assisted selection in aquaculture. 2014;15:376–96.
 54. Wang CM, Lo LC, Zhu ZY, Yue GH. A genome scan for quantitative trait loci affecting growth-related traits in an F1 family of Asian seabass (*Lates calcarifer*). *BMC Genomics;* 2006.
 55. Wang CM, Lo LC, Feng F, Zhu ZY, Yue GH. Identification and verification of QTL associated with growth traits in two genetic backgrounds of Barramundi (*Lates calcarifer*). *Anim Genet.* 2008;39(1):34-39.
 56. Houston RD, Bishop SC, Hamilton A, Guy DR, Tinch AE, Taggart JB, Derayat A, McAndrew BJ, Haley CS. Detection of QTL affecting harvest traits in a commercial Atlantic salmon population. *Anim Genet.* 2009;40(5):753-755.
 57. Houston RD, Haley CS, Hamilton A, Guy DR, Tinch AE, Taggart JB. Major quantitative trait loci affect resistance to infectious pancreatic necrosis in Atlantic salmon (*Salmo salar*). *Genetics.* 2009;178:1109–15.
 58. Song W, Li Y, Zhao Y, Liu Y, Niu Y, Pang R. Construction of a high density microsatellite genetic linkage map and mapping of sexual and growth-related traits in half-smooth tongue sole (*Cynoglossus semilaevis*). *PLoS One.* 2012;7:e52097.
 59. Song W, Pang R, Niu Y, Gao F, Zhao Y, Zhang J. Construction of high density genetic linkage maps and mapping of growth-related quantitative trait loci in the Japanese flounder (*Paralichthys olivaceus*). *PLoS One.* 2012;7:e50404; 2012
 60. Li J, Boroevich KA, Koop BF, Davidson WS. Comparative genomics identifies candidate genes for Infectious Salmon Anemia (ISA) resistance in Atlantic Salmon (*Salmo salar*). *Mar Biotechnol.* 2009;13:232-241.
 61. Wang S, Meyer E, McKay JK, Matz MV. 2b-RAD: a simple and flexible method for genome-wide genotyping. *Nat Methods.* 2012;9:808–10.
 62. Davidson WS, Koop BF, Jones SJM, Iturra P, Vidal R, Mas A, Jonassen I, Lien S, Omholt SW. Sequencing the genome of the Atlantic salmon (*Salmo salar*). *Genome Biol.* 2009;11:403.
 63. Eshel O, Shirak A, Weller JI, Slossman T, Hulata G, Cnaani A. Fine mapping of a locus on linkage group 23 for sex determination in Nile tilapia (*Oreochromis niloticus*). *Anim Genet.* 2011;42:222–4.
 64. Moghadam H, Poissant J, Fotherby H, Haidle L, Ferguson M, Danzmann R. Quantitative trait loci for body weight, condition factor and age at sexual maturation in Arctic charr (*Salvelinus*

- alpinus*): comparative analysis with rainbow trout (*Oncorhynchus mykiss*) and Atlantic salmon (*Salmo salar*). Mol Genet Genomics. 2007;277(6):647-661.
65. Gharbi K, Gautier A, Danzmann RG, Gharbi S, Sakamoto T, Hoyheim B, Taggart JB, Cairney M, Powell R, Krieg F, Okamoto N, Ferguson MM, Holm LE, Guyomard R. A linkage map for brown trout (*Salmo trutta*): Chromosome homeologies and comparative genome organization with other salmonid fish. Genetics. 2006;172:2405–2419.
66. Gilbey J, Verspoor E, McLay A, Houlihan D. A microsatellite linkage map for Atlantic salmon (*Salmo salar*). Anim Genet. 2004;35:98–105.
67. Woram RA, McGowan C, Stout JA, Gharbi K, Ferguson MM, Hoyheim B, Sakamoto J, Davidson W, Rexroad C, Danzmann RG. A genetic linkage map for Arctic char (*Salvelinus alpinus*): Evidence for higher recombination rates and segregation distortion in hybrid versus pure strain mapping parents. Genome. 2003;47:304–315.
68. Loukovitis D, Sarropoulou E, Tsigenopoulos CS, Batargias C, Magoulas A, Apostolidis AP, Chatziplis D, Kotoulas G. Quantitative trait loci involved in sex determination and body growth in the gilthead sea bream (*Sparus aurata* L.) through targeted genome scan. PLoS ONE. 2011; 6:e16599.
69. Martinez P, Bouza C, Hermida M, Fernandez J, Toro MA, Vera M, Pardo B, Millan A, Fernandez C, Vilas R. Identification of the major sex - Determining Region of Turbot (*Scophthalmus maximus*). Genetics. 2009;183(4):1443-1452.
70. Chen J, Wang Y, Yue Y, Xia X, Du Q, Chang Z. A novel male-specific DNA sequence in the common carp, *Cyprinus carpio*. Mol Cell Probes. 2014;23:235–9.

© 2019 Eze; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50277>