



# Activity of Peroxidase and Catalase Enzymes in Thermotolerant Silkworms *Bombyx mori*. L. in Response to *Beauveria bassiana* (Bals. - Criv.) Vuill. Infection

G. H. Shruthi <sup>a</sup>, Manjunath Gowda <sup>a\*</sup>,  
K. C. Narayanaswamy <sup>a</sup> and P. S. Benherlal <sup>b</sup>

<sup>a</sup> Department of Sericulture, College of Agriculture, UAS, GKVK, Bengaluru-560065, India.

<sup>b</sup> Department of Plant Biotechnology, College of Agriculture, UAS, GKVK, Bengaluru-560065, India.

## Authors' contributions

This work was carried out in collaboration among all authors. Authors MG, KCN and PSB conceptualized the research and designed of the experiments. Author MG contributed the experimental materials. Author GHS executed the experiments and collected data. Authors GHS and MG have carried out data analysis and interpretation. Authors GHS and MG prepared the manuscript. All authors have read and approved the final manuscript.

## Article Information

DOI: <https://doi.org/10.9734/jabb/2024/v27i91292>

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/121638>

Original Research Article

Received: 18/06/2024

Accepted: 20/08/2024

Published: 24/08/2024

## ABSTRACT

With an intention to understand the mechanism of dual stress tolerance, a study on antioxidant enzymes activity in thermotolerant bivoltine silkworm breeds and hybrids under *Beauveria bassiana* infection was conducted at the Department of Sericulture, UAS, GKVK, Bengaluru during 2021-22. Thermotolerant bivoltine breeds viz., B1, B4 and B8 and their hybrids with CSR4 along with Pure

\*Corresponding author: E-mail: [mgowda\\_uas@rediffmail.com](mailto:mgowda_uas@rediffmail.com), [mgowda\\_uas@rediffmail.com](mailto:mgowda_uas@rediffmail.com);

Mysore as a muscardine resistant and CSR4 as a muscardine susceptible breeds were used in this study. All the breeds and hybrids were raised following CRD experimental design and were inoculated with *B. bassiana* immediately after 4<sup>th</sup> moult and another group was reared under normal condition. The haemolymph was collected from these breeds and hybrids at 24, 48, 72, 96 and 120 hours post inoculation (hpi) from both the batches and utilized for the estimation of total protein content and peroxidase and catalase enzyme activities in the haemolymph. The total protein content increased from 24 h to 120 h in all the breeds and hybrids under both control and muscardine inoculation, but being lesser under inoculation. The peroxidase activity was found higher in B1 breed from 48 hpi to 120 hpi under both control and inoculated conditions, but decreased at 24 hpi under inoculation. Catalase activity was enhanced at all time intervals under inoculated conditions in Pure Mysore, B1, B4 and B8 breeds and B4×CSR4 hybrid compared to control condition. Thus increased catalase activity in B1 and B4 breeds and decreased peroxidase activity in B1 breed could be associated with their better performance for survival and economic parameters under inoculated condition.

**Keywords:** Silkworm; thermotolerant; antioxidant; muscardine and inoculation.

## 1. INTRODUCTION

An economically significant poikilothermic insect that produces silk is the mulberry silkworm, *Bombyx mori* L. The natural growth and development of silkworms are greatly influenced by environmental conditions, particularly temperature, which can be raised most effectively between 25 and 28 °C. Any difference from the ambient temperature required for proper physiological and metabolic functions is frequently regarded as thermal stress, which is linked to a variety of physiological stress enhances reactive oxygen species (ROS) production, which in turn causes oxidative damage. Silkworms, like all insects, have enzymatic and non-enzymatic antioxidative defence systems to prevent oxidative damage. Silkworm has developed efficient innate immune system to fight against microbial pathogens. The innate immune system plays a crucial biological role in the limiting the microbial infections by using different immune strategies such as production of antimicrobial peptides (AMPs), generation of reactive oxygen species and formation of melanin Abbas et al. [1]. Production of reactive oxygen species under stress and their quenching is a response to get rid of the stress faced by the insect. Superoxide dismutases (SOD), catalases (CAT), peroxidases (POX), and glutathione S-transferases (GST) are the most important antioxidant enzymes secreted by silkworms Pooja et al. [2].

The superoxide dismutase (SOD) upon infection with bacteria after an initial burst (10 min), sharply declined at 30 min and then significantly enhanced in larval haemolymph, implying the modulation of superoxide anion generation is an

early response as part of the defensive process against pathogen invasion Krishnan et al. [3]. Similarly, Catalase activity in the fat body revealed a positive correlation between the control (28±1<sup>o</sup>C) and high temperature (40±1<sup>o</sup>C) Nabizadeh and Jagadeesh Kumar, [4]. The lipid peroxidation (LPO) levels increased significantly in a time-dependent manner under thermal stress in oriental fruit fly, *Bactrocera dorsalis* (Hendel) Jia et al., [5]. Increase in hydrogen peroxide generation with concomitant increase in LPO and protein carbonyl levels in midgut and hemocytes was observed in *B. mori* exposed to low temperature, hypoxia and NPV infection indicating a possible transitory defense mechanism to avoid oxidative stress induced cell damage. Silkworms are known to respond to fungal infection through production of serine protease inhibitors Li et al. [6] and prophenoloxidase – phenoloxidase cascade Zhang et al. [7].

The studies on stress induced ROS production in silkworm under *B. bassiana* infection and its controlling mechanism is scanty, though it is known that catalase activity is affected under *B. bassiana* infection Rajitha and Savithri, [8].

Earlier studies revealed that thermotolerant bivoltine breeds to *B. bassiana* infection that B1, B2, B4 and B8 to be relatively resistant to fungal infection Keerthana, [9]. To identify productive thermotolerant breeds resistant to *B. bassiana* infection in order to identify dual stress tolerant breeds, the study was intended to further understand the mechanism of defending ROS accumulating in thermotolerant bivoltine silkworm under *B. bassiana* infection.

## 2. MATERIALS AND METHODS

### 2.1 Silkworm Breeds Used

Five silkworm breeds viz., B1, B4, B8, CSR4 and Pure Mysore were procured from Central Sericultural Research and Training Institute, Mysore. Three hybrids B1×CSR4, B4×CSR4 and B8×CSR4 were developed by crossing at Sericulture department, GKVK, Bangalore. These breeds and hybrids were raised by following recommend rearing practices Dandin and Giridar, [10]. The fifth instar silkworms were topically inoculated with *B. bassiana* spores (LC<sub>50</sub>, 9.04×10<sup>4</sup> spores/ml @ 0.5 ml per worm).

### 2.2 Collection and Storage of Haemolymph

The haemolymph was collected after 24h, 48h, 72h, 96h and 120h post inoculation i.e., on the first, second, third, fourth and fifth day of fifth instar in each treatment. The haemolymph was collected from randomly selected fifth instar larvae of each set by cutting the third pair of prolegs. The haemolymph, thus coming out were collected and stored in pre-cooled Eppendorf's tube containing a few crystals of phenylthiourea to prevent oxidation. The samples were labelled and then preserved in deep freezer at -20<sup>o</sup> C until further analysis. The samples were centrifuged at 3000 rpm for 15 minutes to separate out the phenylthiourea crystals and haemocytes. The supernatant was used for the estimation after proper dilution Mahesha et al. [11].

### 2.3 Quantitative Estimation of Protein and Enzyme Activity in Haemolymph

The total proteins in the haemolymph was estimated by following Lowry's method (Lowry et al. [12] using crystalline Bovine Serum Albumin (BSA) as standard. Catalase enzyme activity was determined by the method of Abei [13] and Peroxidase enzyme activity was determined according to Sadashivam and Manickam, [14].

The data were analysed using Completely Randomized Design Sundarraj et al. [15]. The mean values of the experiments were compared by using Duncan's Multiple Range Test (DMRT) Duncan, [16] and presented below.

## 3. RESULTS AND DISCUSSION

### 3.1 Total Protein Content (mg/ml)

In the present study the total protein content was estimated at 24h, 48h, 72h, 96h and 120h time

intervals after post inoculation. Total protein content in haemolymph was more under control compared to infected and it was minimum at 24 hpi and maximum at 120 hpi in all breeds and hybrids. It was observed that protein content to be more in hybrids, followed by thermotolerant bivoltine breeds and it was found to be least in Pure Mysore breed. B8 breed among the breeds recorded maximum total protein content of 243.47 mg/ml at 24 hpi under control and 209.61 mg/ml under inoculated conditions. At 48 hpi B4 breed (280.25 mg/ml) recorded maximum protein content among the breeds under control whereas, under inoculated conditions B8 breed (246.98 mg/ml) recorded maximum protein. From 72 hpi to 120 hpi B4 breed (327.24, 357.60 and 435.85 mg/ml under control, 292.22, 340.09 and 356.14 mg/ml under inoculated condition at 72, 96 and 120 hpi, respectively) recorded maximum total protein under both control and inoculated conditions. Among the hybrids B4×CSR4 hybrid (351.47, 417.15, 455.67, 492.16 and 527.48 mg/ml; 310.61, 368.11, 402.55, 467.93 and 490.41 mg/ml under control and infected conditions, respectively) (Table 1) recorded maximum total protein content at all time intervals from 24 hpi to 120 hpi under both control and inoculated conditions.

Ramesh Babu et al. [17] suggested that the survival ability of an animal to stress majorly depends on its protein synthetic potential, any stress on an animal invokes compensatory metabolic adjustments in its tissues through modifications of proteins. Malik and Malik [18] reported that in the normal larva of *B. mori*, haemolymph protein content showed relatively constant increase from first to fourth stage. In the fifth instar larvae, it increased nearly two-fold on the day four and reached a maximum on day nine.

Among ten thermotolerant studied under muscardine infection, the total protein content in haemolymph was found to decrease from 24 hpi to 72 hpi and again decreased 120 hpi in control, whereas in *B. bassiana* inoculation, it decreased from 24 hpi to 72 hpi and then increased at 120 hpi. The protein content under the stress conditions were much lesser than that of corresponding control Sreejith, [19]. In the present study the same trend was seen among all the breeds and hybrids where their was an increase in total protein content from 24 hpi to 120 hpi. The decrease in total protein content due to the infected could be that the pathogen may have caused the reduction in metabolic activity in the silkworm thus reduced protein

synthesis. This would affect the cocoon production of silk content in the infected silkworm batches. These results are in agreement to the findings of Rajitha and Savithri, [20] in cross breed silkworms.

### 3.2 Peroxidase Enzyme Activity ( $\mu\text{mole/l}$ )

In all the breeds and the hybrids, the peroxidase enzyme activity was found higher in control compared to that of infected ones. Maximum peroxidase activity was observed at 24 hpi in all breeds and hybrids. Minimum enzyme activity was seen at 120 hpi. Among breeds, B1 breed 152.95, 150.23, 144.78, 72.47 and 59.73  $\mu\text{mole/l}$ , 78.73, 75.99, 67.70, 61.02 and 59.28  $\mu\text{mole/l}$  (Fig. 1) has shown highest enzyme activity from 24 hpi to 120 hpi under both control and infected conditions, respectively. Among the hybrids, B1 $\times$ CSR4 and B8 $\times$ CSR4 hybrids have shown higher enzyme activity at all time intervals from 24 hpi to 120 hpi.

Reddy and Venkatappa [21] studied the effect of *Staphylococcus aureus* infection in fifth instar silkworm and found gradual decrease in glutathione peroxidase (GPx) activity among infected group, compared to control. The lowest activity was recorded for GPx (0.121 nmol NADPH/min/mg protein) at 24 hours post infection.

### 3.3 Catalase Enzyme Activity ( $\mu\text{moles/min/ml}$ )

In Pure Mysore breed initially catalase enzyme activity was higher under inoculated condition at 24 hpi and 48 hpi, then it decreased at 72 hpi and increased at 96 hpi up to 120 hpi compared to control. In CSR4 breed the enzyme activity was low at 24 hpi and also at 48 hpi, then it has showed higher activity at 72 hpi, 96 hpi and 120 hpi compared to that of control.

In B1 breed the enzyme activity was higher compared to control conditions throughout the larval period from 24 hpi to 120 hpi compared to control. In B4 breed initially at 24 hpi and 48 hpi enzyme activity was higher under inoculated conditions then it decreased at 72 hpi and again it increased at 96 hpi and 120 hpi compared to control. In B8 breed at 24 hpi enzyme activity was same under both control and inoculated conditions, then it decreased under inoculated

conditions at 48 hpi, increased at 72 hpi, decreased at 96 hpi and again increased at 120 hpi compared to control. In B1 $\times$ CSR4 hybrid initially enzyme activity was same at 24 hpi under both control and inoculated conditions. Then from 48 hpi to 120 hpi enzyme activity was lower under inoculated condition compared to control. In B4 $\times$ CSR4 hybrid the catalase enzyme activity was lower under inoculated conditions up to 48 hpi compared to that of control. Then from 72 hpi to 120 hpi enzyme activity was higher under inoculated conditions than that of controlled ones. B8 $\times$ CSR4 hybrid showed lower enzyme activity lower at 24 hpi under inoculated conditions compared to control, then at 48 hpi enzyme activity increased under inoculated conditions and again from 72 hpi to 120 hpi it decreased under infection compare to control condition (Fig. 2).

Catalase enzyme activity was found to be enhanced under inoculated conditions in the breeds Pure Mysore, B1, B4, B8 and hybrid B4 $\times$ CSR4 compared to control conditions which may be associated with their resistance to the disease. But in CSR4 breed and B1 $\times$ CSR4 and B8 $\times$ CSR4 hybrids enzyme activity was maximum under control compared to inoculated conditions. At different time intervals the breeds and hybrids exhibited more fluctuations in catalase enzyme activity, except in B1 breed and B1 $\times$ CSR4 hybrid which showed higher catalase activity under inoculated and normal conditions, respectively.

According to Praveena and Savithri [22] catalase activity during fifth instar under *Beauveria bassiana* infection in three breeds revealed that the activity gradually decreased in all breeds under infected condition compared to control. Elevated antioxidant enzyme catalase activity in bivoltine double hybrid suggests their higher level of immunity compared to single hybrids. Such enhanced antioxidant enzyme activity may provide defense to the host organism.

The average catalase and peroxidase enzyme activity over the breeds at different hpi is represented in the Fig. 3a and 3b, respectively. These figures indicate that catalase enzyme activity among the breeds under fungal infection was higher compared to that of control condition at 24, 48, 72 and 120 hpi except at 96 hpi. While, the peroxidase enzyme activity was higher in healthy silkworms than under *B. bassiana* infected condition. Hence, it can be presumed

**Table 1. Total protein content (mg/mL) in haemolymph of selected thermotolerant bivoltine silkworm breeds and hybrids as influenced by *B. bassiana* infection**

Breeds and hybrids		24 hpi		48 hpi		72 hpi		96 hpi		120 hpi	
		Control	Muscardine inoculation	Control	Muscardine inoculation	Control	Muscardine inoculation	Control	Muscardine inoculation	Control	Muscardine inoculation
Breeds	Pure mysore	179.26 <sup>g</sup>	160.58 <sup>de</sup>	195.91 <sup>ef</sup>	171.09 <sup>f</sup>	210.20 <sup>f</sup>	172.82 <sup>f</sup>	228.00 <sup>f</sup>	212.24 <sup>e</sup>	266.82 <sup>f</sup>	260.70 <sup>f</sup>
	CSR4	160.58 <sup>h</sup>	150.95 <sup>e</sup>	207.28 <sup>e</sup>	180.72 <sup>f</sup>	237.64 <sup>e</sup>	187.14 <sup>ef</sup>	251.65 <sup>e</sup>	201.73 <sup>e</sup>	329.87 <sup>d</sup>	258.94 <sup>f</sup>
	B1	191.52 <sup>f</sup>	160.29 <sup>de</sup>	192.69 <sup>f</sup>	170.21 <sup>f</sup>	287.55 <sup>d</sup>	214.87 <sup>d</sup>	317.03 <sup>d</sup>	297.18 <sup>d</sup>	311.19 <sup>e</sup>	298.06 <sup>e</sup>
	B4	225.38 <sup>e</sup>	171.96 <sup>d</sup>	280.25 <sup>d</sup>	211.36 <sup>e</sup>	327.24 <sup>c</sup>	292.22 <sup>c</sup>	357.60 <sup>c</sup>	340.09 <sup>c</sup>	435.85 <sup>c</sup>	356.14 <sup>d</sup>
	B8	243.47 <sup>d</sup>	209.61 <sup>c</sup>	268.56 <sup>d</sup>	246.98 <sup>d</sup>	283.46 <sup>d</sup>	196.77 <sup>e</sup>	322.87 <sup>d</sup>	206.99 <sup>e</sup>	318.49 <sup>de</sup>	214.29 <sup>g</sup>
Hybrids	B1xCSR4	263.03 <sup>c</sup>	253.40 <sup>b</sup>	313.82 <sup>c</sup>	271.78 <sup>c</sup>	354.39 <sup>b</sup>	305.35 <sup>c</sup>	430.86 <sup>b</sup>	336.00 <sup>c</sup>	505.00 <sup>b</sup>	434.95 <sup>b</sup>
	B4xCSR4	351.47 <sup>a</sup>	310.61 <sup>a</sup>	417.15 <sup>a</sup>	368.11 <sup>a</sup>	455.67 <sup>a</sup>	402.55 <sup>a</sup>	492.16 <sup>a</sup>	467.93 <sup>a</sup>	527.48 <sup>a</sup>	490.41 <sup>a</sup>
	B8xCSR4	338.63 <sup>b</sup>	243.77 <sup>b</sup>	348.55 <sup>b</sup>	315.57 <sup>b</sup>	345.92 <sup>b</sup>	329.87 <sup>b</sup>	440.79 <sup>b</sup>	358.77 <sup>b</sup>	450.13 <sup>c</sup>	385.91 <sup>c</sup>
F test	*	*	*	*	*	*	*	*	*	*	*
S.Em ±	3.733	4.126	4.115	4.026	4.035	4.869	4.774	5.773	5.065	4.676	
CD @5%	11.287	12.475	12.442	12.174	12.2	14.724	14.435	17.455	15.314	14.14	
CV (%)	2.648	3.441	2.563	2.882	2.234	3.21	2.328	3.304	2.231	2.4	

Figures with same superscript are statistically on par; \*-Significant @5%






Fig. 1. Peroxidase enzyme activity in thermotolerant bivoltine silkworm breeds and hybrids under control and muscardine inoculation at different time intervals



**Fig. 2. Catalase enzyme activity in thermotolerant bivoltine silkworm breeds and hybrids under control and muscardine inoculation at different time intervals**



**Fig. 3. Enzyme activity of the breeds over different time intervals (a) catalase and (b) peroxidase. Each value is derived from 8 breeds/hybrids at 24, 48, 72, 96 & 120 hpi with *B. bassiana* @  $9.04 \times 10^4$  spores / ml on the first day of 5<sup>th</sup> instar.**

that the breeds tried to respond to infection by increasing catalase activity rather than enhancing peroxidase activity.

Catalase activity increased in the larvae of *Spodoptera litura* as a response to *Aspergillus flavus* infection in accordance with increase in load of the fungus Karthi et al. [23]. Similar phenomenon is observed in the present study where catalase activity increased over a period of 5 days among thermotolerant bivoltine silkworm breeds in response to *B. bassiana* infection. In the fruit fly, *Bactrocera dorsalis*, the peroxidase activity decreased in response to temperature stress, either lower or higher temperature than 27 °C, regardless of duration of exposure to thermal stress Jia et al. [5].

#### 4. CONCLUSION

This study investigated the response of antioxidant enzyme activity in thermotolerant bivoltine silkworm breeds and hybrids under *Beauveria bassiana* infection. Key findings revealed significant variations in total protein content, peroxidase and catalase enzyme activities among different breeds and hybrids under both control and infected conditions. The study suggests that the observed increase in catalase activity, particularly in B1 and B4 breeds, along with the decrease in peroxidase activity in B1, could serve as biochemical markers for selecting better performing silkworm breeds under fungal infection. These findings provide valuable insights into the mechanisms of

dual stress tolerance in silkworms and could aid in the selection and development of more resilient silkworm breeds for sericulture.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of manuscripts.

#### ACKNOWLEDGEMENT

The authors are highly grateful to CSRTI, Mysore, for supplying thermotolerant bivoltine silkworm breeds for the study. First author is thankful to department of plant biotechnology, UAS, GKVK, Bengaluru-65 for providing lab facilities for enzyme assay.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

1. Abbas MN, Kausar S, Zhao Y, Cui H. Immune strategies of silkworm, *Bombyx mori* against microbial infections. *Invertebr. Surviv. J.* 2019;16:130-140.
2. Pooja M, Rahul K, Chattopadhyay S, Sivaprasad V. Effect of thermal stress on antioxidant responses in *Bombyx mori*. *Chem. Rev. Lett.* 2021;10(38):288-294.



3. Krishnan N, Chattopadhyay S, Kundu JK, Chaudhuri A. Superoxide dismutase activity in haemocytes and haemolymph of *Bombyx mori* following bacterial infection. *Curr. Sci.* 2002;83(3):321-325.
4. Nabizadeh P, Jagadeesh Kumar TS. Fat body catalase activity as a biochemical index for the recognition of thermotolerant breeds of mulberry silkworm *Bombyx mori* L. *J. Therm. Biol.*, 2010;36:1-6.
5. Jia FX, Dou W, Hu F, Wang JJ. Effects of thermal stress on lipid peroxidation and antioxidant enzyme activities of oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *Fla. Entomol.* 2011;94(4): 956-963.
6. Li Y, Zhao P, Liu S, Dong Z, Chen J, Xiang Z, Xia Q. A novel protease inhibitor in *Bombyx mori* is involved in defense against *Beauveria bassiana*. *Insect Biochem. Mol. Biol.* 2012;42:766-775.
7. Zhang J, Huang W, Yuan C, Lu Y, Yang B, Wang CY, Zhang P, Dobens L, Zou Z, Wang C, Ling E. Prophenoloxidase-mediated Ex vivo immunity to delay fungal infection after insect ecdysis. *Front. Immunol.* 2017;8:1-14.
8. Rajitha K, Savithri G. Dynamics of catalase in the haemolymph of silkworm *Bombyx mori* L. during the progress of fungal pathogen *Beauveria bassiana* (Bals.) Vuill. *IJBPSA.* 2014;5(1):62-65.
9. Keerthana A. Studies on response of thermotolerant bivoltine silkworm breeds to white muscardine infection. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Bangalore, Indi. 2018;1-96.
10. Dandin SB, Giridhar K. Handbook of sericulture technologies, CSB publications. 2010;13.
11. Mahesha HB, Thejaswini PH, Honnaiah S. Haemolymph proteins of F1 progeny raised from ethyl methane sulfonate treated silkworm *Bombyx mori* L. *Ind J Seri.* 2000; 39 (2):139-144.
12. Lowry OH, Rosebrough NJ, Farr AL, Randal RJ. Protein measurement with the Folin phenol reagent. *J. Biochem.* 1951; 193:265-275.
13. Aebi H. Catalase In-vitro. *Methods Enzymol.* 1984;105:121-126.
14. Sadasivam S, Manickam A. Biochemical methods. New Age International Publishers, New Delhi. 2004;108-109.
15. Sundararaj N, Nagaraju S, Venkataramu MN, Jagannath MK. Designs of analysis of field experiments. UAS, Bangalore. 1972; 424.
16. Duncan F. Multiple range test and multiple 'F' test. *Biometrics.* 1955;11:1-42.
17. Ramesh Babu K, Ramakrishna S, Harish Kumar Reddy Y, Lakshmi G, Naidu NV, Basha SS, Bhaskar M. Metabolic alterations and Molecular mechanism in silkworm larvae during viral infection: A review. *Afr. J. Biotechnol.* 2009;8(6):899-907.
18. Malik MA, Malik FA. Effect of high temperature on protein and free amino acid levels in three selected silkworm races. *Uttar Pradesh Journal of Zoology.* 2009;29:263-268.
19. Sreejith V. Biochemical changes in thermotolerant bivoltine breeds during high temperature treatment and muscardine disease. M.Sc. (Agri.) Thesis, University of Agricultural Sciences, Bangalore, India. 2019;1-154.
20. Rajitha K, Savithri G. Electrophoretic banding pattern of haemolymph proteins of silkworm *Bombyx mori* L. infected with fungal pathogen *Beauveria bassiana* (Bals.) Vuill. *Int. J. Adv. Lif. Sci.* 2013;6(5):447-449.
21. Reddy HA, Venkatappa B. Effect of *Staphylococcus aureus* infection on biochemical and antioxidant activities of fifth instar silkworm larvae (*Bombyx mori* L.). *J Bacteriol Parasitol.* 2016;7(4): 286.
22. Praveena SM, Savithri G. Dynamics of immune enzymes in the haemolymph of different breeds of silkworm *Bombyx mori* L during the progress of fungal pathogen *Beauveria bassiana*. *J. Crit. Rev.* 2020; 7(13).

23. Karthi S, Vaideki K, Shivakumar MS, Ponsankar A, Thanigaivel A, Chellappandian M, Srinivasan PV, Pandian CKM, Hunter WB, Nathana SS. Effect of *Aspergillus flavus* on the mortality and activity of antioxidant enzymes of *Spodoptera litura* Fab. (Lepidoptera: Noctuidae) larvae. Pestic Biochem Phys. 2018;149:54-60.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

<https://www.sdiarticle5.com/review-history/121638>